arXiv:0904.4102v1 [math.FA] 27 Apr 2009

APPROXIMATE IDENTITY AND ARENS REGULARITY OF SOME BANACH ALGEBRAS

KAZEM HAGHNEJAD AZAR AND ABDOLHAMID RIAZI

ABSTRACT. Let A be a Banach algebra with the second dual A^{**}. If A has a bounded approximate identity $(=BAI)$, then A^{**} is unital if and only if A^{**} has a *weak* bounded approximate identity*$(=W^*BAI)$. If A is Arens regular and A has a BAI, then A^* factors on both sides. In this paper we introduce new concepts LW^*W and RW^*W- property and we show that under certain conditions if A has LW^*W and RW^*W- property, then A is Arens regular and also if A is Arens regular, then A has LW^*W and RW^*W- property. We also offer some applications of these new concepts for the special algebras $l^1(G)$, $L^1(G)$, $M(G)$, and $A(G)$.

Mathematics Subject Classification: 46L06; 46L07; 46L10; 47L25
Keywords: Approximate identity, Arens regularity, LW^*W- property, Topological center, Dual space.

Preliminaries

Through of this paper, A is a Banach algebra and A^*, A^{**}, respectively, are the first and second dual of A. We say that a bounded net $(e_\alpha)_{\alpha\in I}$ in A is an approximate identity $(=BAI)$ if, for each $a\in A$, $ae_\alpha\longrightarrow a$ and $e_\alpha a\longrightarrow a$. For $a\in A$ and $f\in A^*$, we denote by $f.a$ and aof respectively, the functionals on A^* defined by $<f.a,b>=<f,ab>=f(ab)$ and $<aof,b>=<f,ba>=f(ba)$. As since the Banach algebra A can be considered a subspace of its second dual, so we can define $<a,f>$ as $<f,a>$ for every $a\in A$ and $f\in A^*$.

We denote by A^*A and AA^*, respectively, $\{f.a:\ a\in A\ and\ f\in A^*\}$ and $\{aof:\ a\in A\ and\ f\in A^*\}$ clearly these two sets are subsets of A^*. We say that the Banach algebra A is unital if there exists an element $e\in A$ such that $e.x=x.e=x$ for each $x\in A$.

Let A be a Banach algebra with a BAI. If the equality $A^*A=A^*$, $(AA^*=A^*)$ holds, then we say that A^* factors on the left (right). If both equalities $A^*A=AA^*=A^*$ hold, then we say that A^* factors on both sides.

Arens [1] has shown that given any Banach algebra A, there exist two algebra multiplications on the second dual of A which extend multiplication on A. In the following, we introduce both multiplication which are given in [11].

Let $a, b \in A$ and $f \in A^*$ and $F, G \in A^{**}$ then the first Arens multiplication is defined as

$$< f.a, b >=< f, ab >,$$

$$< F.f, a >=< F, f.a >,$$

$$< F.G, f >=< F, G.f > .$$

Clearly $F.f \in A^*$ and $F.G \in A^{**}$. We use the notions $(A^{**}, .)$ for A^{**} equipped with first Arens multiplication also we use fa instead of $f.a$ for all $a \in A$ and $f \in A^*$.
The second Arens product is defined as follows
For $a, b \in A$, $f \in A^*$ and $F, G \in A^{**}$ the elements aof , foF of A^* and FoG of A^{**} are defined respectively by the equalities

$$< aof, b >=< f, ba >,$$

$$< foF, a >=< F, aof >,$$

$$< FoG, f >=< G, foF > .$$

An element E of A^{**} is said to be a mixed unit if E is a right unit for the first Arens multiplication and a left unit for the second Arens multiplication. That is, E is a mixed unit if and only if, for each $F \in A^{**}$, $F.E = EoF = F$. We say that A^{**} is unital with respect to the first Arens product, if there exists an element $E \in A^{**}$ such that $F.E = E.F = F$ for all $F \in A^{**}$, and A^{**} is unital with respect to the second Arens product , if there exists an element $E \in A^{**}$ such that $FoE = EoF = F$ for all $F \in A^{**}$. By [3, p.146], an element E of A^{**} is mixed unit if and only if it is a $weak^*$ cluster point of some BAI $(e_\alpha)_{\alpha \in I}$ in A.
We say that A^{**} has a $weak^*$ $bounded\ left\ approximate\ identity (= W^*BLAI)$ with respect to the first Arens product, if there is a bounded net as $(e_\alpha)_\alpha \subseteq A$ such that for all $F \in A^{**}$ and $f \in A^*$, we have $< e_\alpha.F, f >\to< F, f >$. The definition of W^*RBAI is similar to W^*LBAI and if A^{**} has both W^*LBAI and W^*RBAI, then we say that A^{**} has W^*BAI.
Suppose that $F, G \in A^{**}$ and $F.G$, FoG are the first and second Arens Multiplications in A^{**}, respectively. Then the mapping $F \to F.G$, for G fixed in A^{**}, is $weak^* - weak^*$ continuous, but the mapping $F \to G.F$ for G fixed in A^{**} is not in general $weak^* - weak^*$ continuous on A^{**} unless $G \in A$.
As an example for the algebras $L^1(G)^{**}$ and $M(G)^{**}$ whenever G is an infinite topological group the mapping $F \to G.F$, in general, is not $weak^* - weak^*$ *continuous on* $L^1(G)^{**}$ and $M(G)^{**}$, see [10,11]. Hence, the first topological center of A^{**} with respect to first Arens product is defined as follows

$$Z_1 = \{G \in A^{**} : \ F \longrightarrow G.F \ is\ w^* - w^* - continuous\ on\ A^{**}\}.$$

Also, for the second Arens multiplication in A^{**} we know that $aof, foF \in A^*$ and $FoG \in A^{**}$. For G fixed in A^{**} , the mapping $F \rightarrow GoF$ is $weak^* - weak^*$ continuous on A^{**}, but the mapping $F \rightarrow FoG$ is not in general $weak^* - weak^*$ continuous on A^{**} unless $G \in A$. Whence, the second topological center of A^{**} with respect to second Arens product is defined as follows

$$Z_2 = \{G \in A^{**} : \; F \longrightarrow FoG \; is \; w^* - w^* - continuous \; on \; A^{**}\}.$$

It is clear that $A \subseteq Z_1 \bigcap Z_2$ and $Z_1, \; Z_2$ are closed subalgebras of A^{**} endowed with the first second Arens multiplication, respectively.

If, for each $F, G \in A^{**}$, the equality $F.G = FoG$ holds, then the algebra A is said to be Arens regular, see [1,2]. In this case $Z_1 = Z_2 = A^{**}$.
The other extreme situation is that $Z_1 = A$, in this case A is called left strongly Arens irregular, see [9,10,14].
We recall that the topological center of A^{**} is defined the set of all functionals $F \in A^{**}$ which satisfy $F.G = FoG$ for all $G \in A^{**}$, see [11]. In other words, the topological centers of A^{**} with respect to the first and second Arens products can also be defined as following sets respectively

$$Z_1 = \{F \in A^{**} : \; F.G = FoG \quad \forall G \in A^{**}\},$$

$$Z_2 = \{F \in A^{**} : \; G.F = GoF \quad \forall G \in A^{**}\}.$$

For Banach algebra A the topological center of the algebra $(A^*A)^*$ is defined to be following set, see [11].

$$\widetilde{Z} = \{\mu \in (A^*A)^* : \; \lambda \rightarrow \; \lambda.\mu \;\; is \; w^* - w^* - continuous \; on \; (A^*A)^*\}.$$

For all $a \in A$, $f \in A^*$ and $F \in A^{**}$ we have the following statements
i) $f.a = foa$ and $a.f = aof$.
ii) $a.F = aoF$ and $F.a = Foa$.
Because if $b \in A$, then we have

$$< foa, b >=< b, foa >=< ab, f >= f(ab).$$

Also, we have $< f.a, b >= f(ab)$, and we conclude that $f.a = foa$.
Consequently we can write the following relations

$$< a.F, f >=< a, F.f >=< F.f, a >=< F, f.a >=< F, foa >=< aoF, f > .$$

Thus, we conclude that $a.F = aoF$.
In above, part (ii) shows that A^{**} has W^*LBAI with respect to the first Arens product if and only if it has W^*LBAI with respect to the second Arens product.

A functional f in A^* is said to be *wap* (weakly almost periodic) on A if the mapping $a \rightarrow f.a$ from A into A^* is weakly compact. Pym in [15] showed that this definition to the equivalent following condition
For any two net $(a_\alpha)_\alpha$ and $(b_\beta)_\beta$ in $\{a \in A : \ \parallel a \parallel \leq 1\}$, we have

$$\lim_\alpha \lim_\beta < f, a_\alpha b_\beta >= \lim_\beta \lim_\alpha < f, a_\alpha b_\beta >,$$

whenever both iterated limits exist. The collection of all *wap* functionals on A is denoted by $wap(A)$. Also we have $wap(A) = A^*$ if and only if $< F.G, f >=< FoG, f >$ for every $F, G \in A^{**}$, see [19].
In this paper, the notations WSC is used for weakly sequentially complete Banach algebra A and WCC, for weakly completely continuous Banach algebra A, that is, A is said to be weakly sequentially complete, if every weakly Cauchy sequence in A has a weak limit, and A is said to be WCC, if for each $a \in A$, the multiplication operator $x \rightarrow ax$ is weakly compact.
In this paper when A^{**} is referred to as an algebra it will be with respect to the first Arens product unless the second is mentioned explicitly.

Main results

Theorem 1. Let A has a BAI $(e_\alpha)_\alpha$. Then the following assertions hold:
i) For all $f \in A^*$, we have $f.e_\alpha \overset{w^*}{\rightarrow} f$ and $e_\alpha o f \overset{w^*}{\rightarrow} f$.
ii) A^* factors on the right if and only if A^{**} has a W^*LBAI.
iii) A^* factors on the left if and only if A^{**} has a W^*RBAI.
iv) If A^* factors on the left and $E \in A^{**}$ such that $e_\alpha \overset{w^*}{\rightarrow} E$, then E is unit element of A^{**}. Also, if A^* factors on the right and $E \in A^{**}$ such that $e_\alpha \overset{w^*}{\rightarrow} E$, then E is unit element of (A^{**}, o).
v) If A^* factors on the left, then $w^* - closure(AA^{**}) = A^{**}$.

Proof. i) The proof of (i) is clear.
ii) Let $A^*A = A^*$, and $F \in A^{**}$, $f \in A^*$. Then, by [11, 2.1] , we have $f.e_\alpha \overset{w}{\rightarrow} f$, and so

$$< e_\alpha.F, f >=< e_\alpha, F.f >=< F.f, e_\alpha >=< F, f.e_\alpha >\longrightarrow< F, f > .$$

Consequently, $e_\alpha.F \overset{w^*}{\rightarrow} F$.
Conversely, let $e_\alpha.F \overset{w^*}{\rightarrow} F$ for all $F \in A^{**}$. Then, for every $f \in A^*$, we have $< e_\alpha.F, f >\longrightarrow< F, f >$. Since $< e_\alpha.F, f >=< F, f.e_\alpha >$, we have
$< F, f.e_\alpha >\longrightarrow< F, f >$. Therefore by [11, 2.1], we are done.
iii) The proof of (iii) is the same as (ii).

iv) Let $e_\alpha \overset{w^*}{\rightarrow} E$. Then, for all $F \in A^{**}$, by using [11, 2.1], we have the following relations:

$$< E.F, f >=< E, F.f >= \lim_\alpha < e_\alpha, F.f >= \lim_\alpha < F, f.e_\alpha > \rightarrow < F, f > .$$

Hence, we conclude that $E.F = F$.
Now let $f \in A^*$ and $a \in A$. Then , we have

$$< E.f, a >=< E, f.a >= \lim_\alpha < e_\alpha, f.a >= \lim_\alpha < f, a.e_\alpha >=< f, a >$$

Consequently, we have $< F.E, f >=< F, E.f >=< F, f >$ hence $F.E = F$ which implies that E is a unit element of A^{**}.
The next assertion is similar.
v) Let $A^*A = A^*$, and let $f \in A^*$. By [11, 2.1], we have $f.e_\alpha \overset{w}{\rightarrow} f$. For each $F \in A^{**}$, since $e_\alpha.F \in AA^{**}$, we have

$$< e_\alpha.F, f >=< F, f.e_\alpha > \rightarrow < F, f > .$$

Thus, we conclude that $e_\alpha.F \overset{w^*}{\rightarrow} F$ and so $F \in w^* - closure(AA^{**})$.

By the above theorem and by [11, 2.2.a], for the Banach algebra A which has BAI, we conclude that A^{**} has W^*LBAI if and only if $\{F \in A^{**} : A.F \subseteq A\} \subseteq Z_1$, and also A^{**} has W^*RBAI if and only if $\{F \in A^{**} : FoA \subseteq A\} \subseteq Z_2$. For $A = L^1(G)$ where G is locally compact finite group, we have $\{F \in A^{**} : A.F \subseteq A\} \subseteq Z_1 = L^1(G)^{**}$ hence $L^1(G)^{**}$ has W^*LBAI. But when G in locally compact infinite group this relation is not satisfied for $L^1(G)^{**}$ consequently $L^1(G)^{**}$ has not W^*LBAI.

Corollary 2. Let A has BAI. Then the following statements hold.
i) The algebra $(A^{**}, .)$ is unital if and only if A^{**} has W^*LBAI.
ii) The algebra (A^{**}, o) is unital if and only if A^{**} has W^*RBAI.
iii) A^{**} and (A^{**}, o) are unital if and only if A^{**} has W^*BAI.
iv) Suppose that A is a WSC with sequentially BAI. Then A^{**} has W^*LBAI if and only if A^{**} has W^*RBAI, and also $(A^{**}, .)$ is unital if and only if (A^{**}, o) is unital.

Proof: Using Theorem 1 and [11, 2.2], the proofs of (i) , (ii) are clear. The proof of (iv), follows from [11, 2.6] and Theorem 1.

As another application of Theorem 1, let Ω be spectrum of $L^\infty([0, 1])$. Since $L^\infty([0, 1])$ dose not factors on the left and right, then by Theorem 1, $L^1([0, 1])^{**}$

has not W^*LBAI and W^*RBAI, and so $M(\Omega)=L^1([0,1])^{**}$ has not W^*LBAI and W^*RBAI. Consequently by Corollary 2, $M(\Omega)$ is not unital which it follows that Ω is not a group. For more details see [9,16].

Corollary 3. Let A be Arens regular with a BAI. Assume also that A is a two sided ideal in A^{**}, then A^{**} has a W^*BAI and so is unital.

Proof. Use Theorem 1 and [17, 3.2].

Theorem 4. Suppose $f \in wap(A)$ and $G \in A^{**}$ such that $a_\alpha \overset{w^*}{\to} G$ where $(a_\alpha)_\alpha \in A$. Then $f.a_\alpha \overset{w}{\to} f.G$.

Proof. First we show that $f.a_\alpha \overset{w^*}{\to} f.G$. Let $b \in A$, so we have

$$< f.a_\alpha, b >=< a_\alpha, b.f >\to< G, b.f >=< f.G, b > .$$

Thus we have $f.a_\alpha \overset{w^*}{\to} f.G$.
Since $f \in wap(A)$, $< G.F, f >=< GoF, f >$ for all $F, G \in A^{**}$. We have

$$< f.G, F >=< G.F, f >=< GoF, f >=< F, foG >=< foG, F >,$$

it follows that $f.G = foG \in A^*$. For every $F \in A^{**}$, there is $(b_\beta)_\beta \subseteq A$ such that $b_\beta \overset{w^*}{\to} F$, so we have

$$\lim_\alpha < F, f.a_\alpha >= \lim_\alpha \lim_\beta < b_\beta, f.a_\alpha, >= \lim_\alpha \lim_\beta < f, a_\alpha b_\beta >,$$

$$= \lim_\beta \lim_\alpha < f, a_\alpha b_\beta >= \lim_\beta < f.G, b_\beta >= \lim_\beta < b_\beta, f.G >,$$

$$=< F, f.G > .$$

Thus, we conclude that $f.a_\alpha \overset{w}{\to} f.G$.

Corollary 5. Suppose A has a BAI $(e_\alpha)_\alpha$ and $f \in wap(A)$. Then $f.e_\alpha \overset{w}{\to} f$.

Proof. By [5,p.146], there is a mixed unit $E \in A^{**}$ such that $e_\alpha \overset{w^*}{\to} E$. Since $f \in wap(A)$, $< E.F, f >=< EoF, f >=< F, f >$ for all $F \in A^{**}$. So, by Theorem 4, we have $f.e_\alpha \overset{w}{\to} f.E = f$.

Corollary 6. Suppose A is Arens regular and A has a BAI. Then we have the following assertions
i) A^* factors on both sides.
ii) A^{**} is unital, and if A is WSC then A is also unital.

Proof. i) Since A is Arens regular, $wap(A) = A^*$ hence by Corollary 5, we have $f.e_\alpha \xrightarrow{w} f$ for all $f \in A^*$. Then, it follows that A^* factors on the left by [11, 2.1]. Since A is Arens regular, by [11, 2.10] A^* factors on the right and so A^* factors on both sides.
ii) By [11, 2.2], since A^* factors on both sides, $(A^{**}, .)$ and (A^{**}, o) are unital. The second claim hold by using [2, 2.6] and part (i).

Since $L^1(G)^*$, $M(G)^*$ and $A(G)^*$ dose not factors on the left and right whenever G is infinite group, by Corollary 6 we obtain that $L^1(G)^{**}$, $M(G)^{**}$ and $A(G)^{**}$ are not Arens regular.

Theorem 7. For Banach algebra A we have the following assertions
i) Assume that $f \in A^*$ and T_f is the linear operator from A into A^* defined by $T_f a = f.a$. Then, $f \in wap(A)$ if and only if the adjoint of T_f is $weak^* - weak$ continuous.
ii) If A has a BAI and $\widetilde{Z} = (A^*A)^*$, then $A^*A \subseteq wap(A)$.

Proof. i) Let $f \in wap(A)$ and $T_f^* : A^{**} \to A^*$ be the adjoint of T_f. Thus for every $F \in A^{**}$ and $a \in A$, we have $< T_f^* F, a >=< F, T_f a >$.
Suppose $(F_\alpha)_\alpha \subseteq A^{**}$ such that $F_\alpha \xrightarrow{w^*} F$. We show that $T_f^* F_\alpha \xrightarrow{w} T_f^* F$. Let $G \in A^{**}$ and $(a_\beta)_\beta \subseteq A$ such that $a_\beta \xrightarrow{w^*} G$. Since $f \in wap(A)$, we have $< G.F_\alpha, f > \to < G.F, f >$. Hence for fixed α we have the following relations

$$< G, T_f^* F_\alpha >= lim_\beta < T_f^* F_\alpha, a_\beta >= lim_\beta < F_\alpha, T a_\beta >= lim_\beta < F_\alpha, f.a_\beta >$$

$$= lim_\beta < F_\alpha.f, a_\beta >=< G, F_\alpha.f >,$$

hence

$$lim_\alpha < G, T_f^* F_\alpha >= lim_\alpha < G, F_\alpha.f >= lim_\alpha < G.F_\alpha, f >= lim_\alpha < G.F, f >$$

$$= lim_\alpha < G, F.f >=< G, T_f^* F > .$$

We conclude that $T_f^* F_\alpha \xrightarrow{w} T_f^* F$, thus T_f^* is $weak^* - weak$ continuous.
Conversely, let T^* be $weak^* - weak$ continuous and let $(F_\alpha)_\alpha \subseteq A^{**}$ be such that $F_\alpha \xrightarrow{w^*} F$. Then we have

$$< G.F_\alpha, f >=< G, T_f^* F_\alpha > \to < G, T_f^* F >=< G.F, f > .$$

It follow that $f \in wap(A)$.
ii) Let $\widetilde{Z} = (A^*A)^*$ and $F \in A^{**}$. We define $\mu \in (A^*A)^*$ to be the restriction of F to $(A^*A)^*$. Since $\widetilde{Z} = (A^*A)^*$, $\mu \in \widetilde{Z}$. Also for all $a \in A$, we have $a\mu = aF$.

By [11, 3.2], since $A\widetilde{Z} = AZ_1$ and $AZ_1 \subseteq Z_1$, we have $aF \in Z_1$. Therefore, for all $G \in A^{**}$ and $f \in A^*$ we have the following relations

$$< F.G, f.a >=< aF.G, f >=< aFoG, f >=< aFoG, f >=< FoG, f.a > .$$

Consequently $f.a \in wap(A)$.

Theorem 7 shows that a Banach algebra A is Arens regular if and only if T_f^* is $weak^* - weak$ continuous for every $f \in A^*$.

Now we define new concepts namely LW^*W-property and RW^*W-property for Banach algebra A and compare them with Arens regularity of Banach algebra A.

Definition 8. Let $(f_\alpha)_{\alpha \in I} \subseteq A^*$. If for $a \in A$, the convergence $a.f_\alpha \overset{w^*}{\rightarrow} 0$ implies $a.f_\alpha \overset{w}{\rightarrow} 0$, then we say that a has $Left - Weak^* - Weak$ property or LW^*W-property with respect to the first Arens product. The definition of RW^*W-property is similar.
We say that A has the LW^*W-property if, for every $a \in A$, a has LW^*W-property.

Remark and Example 9. We know that if A is reflexive, then $weak$ and $weak^*$ topologies coincide consequently A has LW^*W and RW^*W-property, so it is clear that if A has a member which has not LW^*W-property or RW^*W-property, then A is not reflexive. If A is commutative, then it is clear that the LW^*W-property and RW^*W-property are equivalent. For Banach algebras $l^1(G)$ and $M(G)$ where G is an infinite group, the unit elements of them have not LW^*W-property or RW^*W-property (We know that $l^\infty(G)^* \neq l^1(G)$ and $M(G)^{**} \neq M(G)$).
For a Banach algebra A, if $A^{**} = A^{**}c$ or $A^* = cA^*$ for some $c \in A$ and c has LW^*W-property, then it is easy to see that $A = A^{**}$, and also if $A^{**} = cA^{**}$ or $A^* = A^*c$ for some $c \in A$ and c has RW^*W-property, then $A = A^{**}$.
Now we give some examples of the Banach algebras that have LW^*W-property or RW^*W-property.
Let S be a compact semigroup and $A = C(S)$. Then every $f \in C(S)$ such that $f \geq \alpha$ where $\alpha \in (0, \infty)$ has LW^*W-property. For prove this, let $(\mu_\alpha)_\alpha \subseteq C(S)^* = M(S)$ and $\mu_\alpha.f \overset{w^*}{\rightarrow} 0$ hence $< \mu_\alpha.f, g > \rightarrow 0$ for all $g \in C(S)$. Consequently we have

$$< \mu_\alpha.f, g >=< \mu_\alpha, f.g >= \int_S fg d\mu_\alpha \rightarrow 0.$$

If we set $g = \frac{1}{f}$, then $\mu_\alpha \to 0$. Now let $F \in C(S)^{**} = M(S)^*$. Then we have

$$< F, \mu_\alpha.f >=< F.\mu_\alpha, f >= \int_S f dF\mu_\alpha \to 0.$$

It follows that $\mu_\alpha.f \overset{w}{\to} 0$.
In Corollary 15, we will show that if the Banach algebra A is Arens regular, then A has both LW^*W-property and RW^*W-property.

In the following, we design some condition for a Banach algebra A with LW^*W-property or RW^*W-property, so such Banach algebras are Arens regular.

Theorem 10. For Banach algebra A the following statements hold.
i) If $A^{**} = cA^{**}$ for some $c \in A$ and c has LW^*W-property, then A is Arens regular.
ii) If $A^{**} = A^{**}c$ for some $c \in A$ and c has RW^*W-property, then A is Arens regular.
iii) If $A^* = A^*c$ and c has LW^*W-property, then $A^{**}c \subseteq Z_1 \cap Z_2$ and $cA^* \subseteq wap(A) \subseteq A^*c$.
iv) If $A^* = cA^*$ and c has RW^*W-property, then $cA^{**} \subseteq Z_1 \cap Z_2$ and $A^*c \subseteq wap(A) \subseteq cA^*$.

Proof. i) Let $F, G \in A^{**}$. We show that the mapping $F \to G.F$ is $w^* - w^* - continuous$. Let $f \in A^*$ and $(F_\alpha)_\alpha \subseteq A^{**}$ be such that $F_\alpha \overset{w^*}{\to} F$. Then for all $a \in A$ we have

$$< F_\alpha.f, a >=< F_\alpha, f.a >\to< F, f.a >=< F.f, a > .$$

Consequently, we have $F_\alpha.f \overset{w^*}{\to} F.f$ for all $f \in A^*$ which implies that $(F_\alpha.f)a \overset{w^*}{\to} (F.f)a$ for all $a \in A$. Since c has LW^*W-property, $(F_\alpha.f)c \overset{w}{\to} (F.f)c$. Thus for all $G \in A^{**}$, we have

$$< cG, F_\alpha.f >=< G, (F_\alpha.f)c >\to< G, (F.f)c >=< cG, F.f > .$$

Since $A^{**} = cA^{**}$, we can replace cG by G, and consequently we have
$< G.F_\alpha, f >\to< G.F, f >$ for all $f \in A^*$. Therefore $G \in Z_1$ so that $Z_1 = A^{**}$.
ii) The proof of (ii) is similar to (i).
iii) We show that the mapping $F \to Gc.F$ is $w^* - w^*$-continuous whenever $F, G \in A^{**}$. Let $(F_\alpha) \subseteq A^{**}$ be a net such that $F_\alpha \overset{w^*}{\to} F$. Then, we have $(F_\alpha.f) \overset{w^*}{\to} (F.f)$ for every $f \in A^*$, hence $(F_\alpha.f - F.f) \overset{w^*}{\to} 0$. Since $(F_\alpha.f - F.f) \in A^* = A^*c$, there is $(g_\alpha)_\alpha \subseteq A^*$ such that $(F_\alpha.f - F.f) = g_\alpha c$. Then we

have $c(g_\alpha c) \overset{w^*}{\to} 0$, it follow that $c(g_\alpha c) \overset{w}{\to} 0$, since c has LW^*W-property. Consequently for all $G \in A^{**}$, we have $< G, c(g_\alpha c) >\to 0$ which implies $< Gc, (F_\alpha.f - F.f) >\to 0$ hence $Gc \in Z_1$ so $A^{**}c \subseteq Z_1$. Similarly $A^{**}c \subseteq Z_2$. Hence we have $A^{**}c \subseteq Z_1 \cap Z_2$.
Now let $F, G \in A^{**}$. Since $Gc \in Z_2$, we have $Fo(Gc) = F.(Gc)$, consequently for all $f \in A^*$, we have the following statements

$$< FoG, cf >=< Fo(Gc), f >=< F.(Gc), f >=< F.G, cf > .$$

Hence we have $cA^* \subseteq wap(A)$.
iv) The proof is similar as (iii).

Let G be an infinite group. Since $M(G)$ is not Arens regular, by Theorem 10, the unit element of $M(G)$ has not LW^*W-property or RW^*W-property. Similarly conclusions hold for the Banach algebra $l^1(G)$.

Theorem 11. Let A and B be Banach algebras and let h be a bounded homomorphism from A onto B. If A has LW^*W-property (resp. RW^*W-property), then B has LW^*W-property (resp. RW^*W-property).

Proof. Since h is continuous, the second adjoint of h, h^{**} is $weak^* - weak$ continuous from A^{**} into B^{**}. Hence we conclude that $h^{**}(F.G) = h^{**}(F).h^{**}(G)$ for all $F, G \in A^{**}$.
Let $(g_\alpha)_\alpha \subseteq B^*$ be such that $b.g_\alpha \overset{w^*}{\to} 0$ where $b \in B$. We define $h^*(g_\alpha) = f_\alpha \in A^*$ and get $h(a) = b$ for some $a \in A$. Then for every $x \in A$, we have

$$< a.f_\alpha, x >=< f_\alpha, x.a >=< h^*(g_\alpha), x.a >=< g_\alpha, h(x.a) >=< g_\alpha, h(x).h(a) >$$

$$< g_\alpha, h(x).b >=< b.g_\alpha, h(x) >\to 0.$$

Thus $a.f_\alpha \overset{w^*}{\to} 0$. Since a has LW^*W-property, $a.f_\alpha \overset{w}{\to} 0$. Now, let $G \in B^{**}$. Since h is surjective by [12, 3.1.22] there is $F \in A^{**}$ such that $h^{**}(F) = G$. Then we have the following assertions

$$< G, b.g_\alpha >=< G.b, g_\alpha >=< h^{**}(F)h(a), g_\alpha >=< h^{**}(F.a), g_\alpha >$$

$$=< (F.a), h^*(g_\alpha) >=< (F.a), f_\alpha >=< F, a.f_\alpha >\to 0$$

Thus we conclude that $b.g_\alpha \overset{w^*}{\to} 0$, therefore B has LW^*W-property.

Theorem 12. For a Banach algebra A the following assertions hold
i) If $A^*A = A^*$ and A has RW^*W-property, then A is Arens regular.
ii) If $AA^* = A^*$ and A has LW^*W-property, then A is Arens regular.

Proof. i) Let $(F_\alpha) \subseteq A^{**}$ be a net such that $F_\alpha \overset{w^*}{\rightarrow} F$. Then, we have $(F_\alpha.f).a \overset{w^*}{\rightarrow} (F.f).a$ for every $a \in A$ and $f \in A^*$, and so $F_\alpha.(f.a) \overset{w^*}{\rightarrow} F.(f.a)$. Since A has LW^*W-property, we conclude that $F_\alpha.(f.a) \overset{w}{\rightarrow} F.(f.a)$. Therefore for all $G \in A^{**}$, $f \in A^*$ and $a \in A$ we have the following relations

$$< G.F_\alpha, f.a >=< G, F_\alpha(f.a) >=< G, (F_\alpha.f).a >\rightarrow< G, (F.f).a >$$

$$=< G.F, f.a > .$$

Since $A^*A = A^*$, the result follows.
ii) The Proof is similar to (i).

By Theorem 11 and Theorem 12(ii), if A, B are Banach algebras and h is a bounded homomorphism from A onto B whenever B factors on the right, then if A has LW^*W -property, B is Arens regular. Also from Theorem 12(i) and [11, 2.6] we conclude that when A is WSC and $AA^* = A^*$ [or A is unital] and A has RW^*W-property then A is Arens regular.

Lemma 13. For a Banach algebra A the following statements hold.
i) For all $f \in A^*$ and $F \in A^{**}$ there is a net $(x_\alpha)_\alpha \subseteq A$ such that $f.x_\alpha \overset{w^*}{\rightarrow} foF$.
ii) A is Arens regular if and only if $f.x_\alpha \overset{w}{\rightarrow} foF$ where $F \in A^{**}$, $f \in A^*$ and $(x_\alpha)_\alpha \subseteq A$ such that $x_\alpha \overset{w^*}{\rightarrow} F$.

Proof. i) Since $w^* - closureA = A^{**}$, for all $F \in A^{**}$, there is a net $(x_\alpha)_\alpha \subseteq A$ such that $x_\alpha \overset{w^*}{\rightarrow} F$. Now, let $a \in A$. Then, we have

$$< foF, a >=< F, aof >= \lim_\alpha < x_\alpha, aof >= \lim_\alpha < aof, x_\alpha >$$

$$= \lim_\alpha < f, x_\alpha.a >= \lim_\alpha < f.x_\alpha, a > .$$

We conclude that $f.x_\alpha \overset{w^*}{\rightarrow} foF$.
ii) Let $f.x_\alpha \overset{w}{\rightarrow} foF$. Then, for all $G \in A^{**}$, we have

$$< F.G, f >=< F, G.f >= \lim_\alpha < G.f, x_\alpha >= \lim_\alpha < G, f.x_\alpha >=< G, foF >$$

$$=< FoG, f > .$$

Hence $F.G = FoG$ for all $F, G \in A^{**}$.
Conversely, let $F.G = FoG$ for all $F, G \in A^{**}$ and let $x_\alpha \overset{w^*}{\rightarrow} F$. Then, for all $f \in A^*$ and $G \in A^{**}$, we have the following relations

$$< G, f.x_\alpha >=< G.f, x_\alpha >\rightarrow< F, G.f >=< F.G, f >=< FoG, f >$$

$$=< G, foF >,$$

which implies that $f.x_\alpha \overset{w}{\to} foF$.

Theorem 14. For a Banach algebra A the following assertions hold.
i) Suppose $\lim_\alpha \lim_\beta < f_\beta, a_\alpha >= \lim_\beta \lim_\alpha < f_\beta, a_\alpha >$ for every $(a_\alpha)_\alpha \subseteq A$ and $(f_\beta)_\beta \subseteq A^*$. Then, if $A^*A = A^*$ or $AA^* = A^*$ then A is Arens regular. Also if $A^{**} = cA^{**}$ or $A^{**} = A^{**}c$ for some $c \in A$, then A is reflexive.

ii) Let A be Arens regular and let $(a_\alpha)_\alpha \subseteq A$ and $(f_\beta)_\beta \subseteq A^*$ such that have taken limits, respectively, in the $weak^*$ and weak topology in A^{**} and A^*. Then for all $F \in A^{**}$, we have

$$\lim_\alpha \lim_\beta < F, f_\beta.a_\alpha >= \lim_\beta \lim_\alpha < F, f_\beta.a_\alpha > .$$

iii) If for some $a \in A$, we have

$$\lim_\alpha \lim_\beta < f_\beta a, a_\alpha >= \lim_\beta \lim_\alpha < f_\beta a, a_\alpha >,$$

then a has RW^*W-property. Also if

$$\lim_\alpha \lim_\beta < af_\beta, a_\alpha >= \lim_\beta \lim_\alpha < af_\beta, a_\alpha >,$$

then a has LW^*W-property.

Proof. i) If we show that A has both LW^*W-property and RW^*W-property then by Theorem 10 and Theorem 12 , we are done.
Suppose that $(f_\beta)_\beta \subseteq A^*$ and $a \in A$ such that $a.f_\beta \overset{w^*}{\to} 0$. Let $F \in A^{**}$. Since $w^* - closureA = A^{**}$, there is $(a_\alpha)_\alpha \subseteq A$ such that $a_\alpha \overset{w^*}{\to} F$. Consequently, we have the following equalities

$$\lim_\beta < F, a.f_\beta >= \lim_\beta < F.a, f_\beta >= \lim_\beta \lim_\alpha < a_\alpha.a, f_\beta >= \lim_\beta \lim_\alpha < f_\beta, a_\alpha.a >$$

$$= \lim_\alpha \lim_\beta < f_\beta, a_\alpha.a >= \lim_\alpha \lim_\beta < a.f_\beta, a_\alpha >= 0.$$

Therefore, we obtain $a.f_\beta \overset{w}{\to} 0$. Similarly A has RW^*W-property.
ii) Suppose that $(a_\alpha)_\alpha \subseteq A$ such that $a_\alpha \overset{w^*}{\to} G$ for some $G \in A^{**}$. Let $(f_\beta)_\beta \subseteq A^*$ such that $f_\beta \overset{w}{\to} f$ where $f \in A^*$. Then by Lemma 13, for fixed β, we have $f_\beta.a_\alpha \overset{w}{\to} f_\beta oG$. So, for all $F \in A^{**}$ we have the following relations

$$\lim_\beta \lim_\alpha < F, f_\beta.a_\alpha >= \lim_\beta < F, f_\beta.oG >= \lim_\beta < GoF, f_\beta >=< GoF, f >$$

$$=< G.F, f >=< G, F.f >= \lim_{\alpha} < F.f, a_\alpha >= \lim_{\alpha} < F, f.a_\alpha >$$

$$= \lim_{\alpha} < a_\alpha.F, f >= \lim_{\alpha}\lim_{\beta} < a_\alpha.F, f_\beta >= \lim_{\alpha}\lim_{\beta} < F, f_\beta.a_\alpha > .$$

iii) Suppose that $(f_\beta)_\beta \subseteq A^*$ and $f_\beta a \overset{w^*}{\rightarrow} 0$. Let $F \in A^{**}$ and $(a_\alpha)_\alpha \subseteq A$ such that $a_\alpha \overset{w^*}{\rightarrow} F$. Then, we have

$$\lim_{\beta} < F, f_\beta a >= \lim_{\beta} < aF, f_\beta >= \lim_{\beta}\lim_{\alpha} < f_\beta, aa_\alpha >$$

$$= \lim_{\alpha}\lim_{\beta} < f_\beta, aa_\alpha >= \lim_{\alpha}\lim_{\beta} < f_\beta a, a_\alpha >= 0$$

Consequently, we have $f_\beta a \overset{w}{\rightarrow} 0$. The prove of the next assertion is the same.

Definition 15. Let A be Banach algebra. We say that A^* strong left (right) factors, if for all $(f_\alpha)_\alpha \subseteq A^*$, there are $(a_\alpha)_\alpha \subseteq A$ and $f \in A^*$ such that $f_\alpha = f.a_\alpha$ $(f_\alpha = a_\alpha.f)$ where $(a_\alpha)_\alpha \subseteq A$ has limit in the $weak^*$ topology in A^{**}. If A^* strong left and right factors, then we say that A^* strong factors.
For a Banach algebra A with a BAI, it is clears that if A^* is strong left (right) factors, then A^* factors on the left (right).

Theorem 16. Let $AA^* \subseteq wapA$. If A^* strong factors on the left (right), then A has LW^*W-property (RW^*W-property).

Proof. We prove that A has LW^*W-property and the proof of RW^*W-property is similar. Suppose that $a \in A$ and $(f_\alpha)_\alpha \subseteq A^*$ such that $a.f_\alpha \overset{w}{\rightarrow} 0$. Since A^* strong factors, there are $(a_\alpha)_\alpha \subseteq A$ and $f \in A^*$ such that $f_\alpha = f.a_\alpha$ $(f_\alpha = a_\alpha.f)$ where $(a_\alpha)_\alpha \subseteq A$ has limit in the $weak^*$ topology in A^{**}. Let $F \in A^{**}$ and $(a_\beta)_\beta \subseteq A$ such that $b_\beta \overset{w^*}{\rightarrow} F$. Then

$$\lim_{\alpha} < F, a.f_\alpha >= \lim_{\alpha}\lim_{\beta} < af_\alpha, b_\beta >= \lim_{\alpha}\lim_{\beta} < af.a_\alpha, b_\beta >$$

$$= \lim_{\alpha}\lim_{\beta} < af, a_\alpha b_\beta >= \lim_{\beta}\lim_{\alpha} < af, a_\alpha b_\beta >= \lim_{\beta}\lim_{\alpha} < af.a_\alpha, b_\beta >$$

$$= \lim_{\beta}\lim_{\alpha} < af_\alpha, b_\beta >= 0.$$

Then $a \in A$ has LW^*W-property and so A has LW^*W-property.

Theorem 17. Suppose that A has a BAI and $AA^{**} = A^{**}$. If A^* is not right factor then there is $a \in A$ such that a has not RW^*W-property.

Proof. Let $(e_\alpha)_\alpha$ be a BAI for A. Then for all $f \in A^*$, we have $f.e_\alpha \overset{w^*}{\rightarrow} f$,

and so for all $a \in A$, we have $(f.e_\alpha).a \overset{w^*}{\to} f.a$. Suppose on contrary that $a \in A$ has RW^*W-property then $(f.e_\alpha).a \overset{w}{\to} f.a$.
Now, for $F \in A^{**}$ we have

$$< a.F, f.e_\alpha >=< F, (f.e_\alpha).a >=< F.f, e_\alpha.a >\to< F.f, a >=< a.F, f >,$$

hence $f.e_\alpha \overset{w}{\to} f$, since $AA^{**} = A^{**}$. By [11, 2.11] we conclude that $A^*A = A^*$ which is a contradiction.

Open problems
1. Suppose that A has LW^*W-property and A is WCC and WSC. Is A reflexive?
2. For a non abelian Banach algebra A, when LW^*W-property and RW^*W-property are equivalent?
3. If A^* factors on the right and A has RW^*W-property or if A^* factors on the left and A has LW^*W-property, is A Arens regular?
4. If A factors on the left and Arens regular, is A has LW^*W-property?
5. If A is Arens irregular (or extremely Arens irregular), dose A has some member with LW^*W-property or RW^*W-property?

References

[1] R.E. Arens, *The adjoint of a bilinear operation*, Amer. Math. Soc. **2** (1951), 839-848.

[2] N. Arikan, *A simple condition ensuring the Arens regularity of bilinear mapping*, Proc. Amir. Math. Soc. **84** (4) (1982), 525-532.

[3] J.W. Baker, J.S. Pym, H. L. Vasudeva, *Totally ordered measure spaces and their - algebras*, Mathematika. **2** (1982), 42-54.

[4] J. Baker, A.T. Lau, J.S. Pym *Module homomorphism and topological centers associated with weakly sequentially compact Banach algebras*, Journal of Functional Analysis. **158** (1998), 186-208.

[5] F.F. Bonsall, J. Duncan, *Complete normed algebras*, Springer-Verlag, Berlin 1973.

[6] N. Dunford, J.T. Schwartz, *Linear operators.I*, Wiley, New york 1958.

[7] E. Hewitt, K.A. Ross, *Abstract harmonic analysis*, Springer, Berlin, Vol I 1963.

[8] E. Hewitt, K.A. Ross, *Abstract harmonic analysis*, Springer, Berlin, Vol II 1970.

[9] N. Isik, J.S. Pym, A. Ulger, *The second dual of the group algebra of compact groups*, J. London. Math. Soc. **35** (2) (1987), 135-148.

[10] A.T. Lau, V. Losert, *On the second Conjugate Algebra of locally compact groups*, J. London Math. Soc. **37** (2)(1988), 464-480.

[11] A.T. Lau, A. Ulger, *Topological center of certain dual algebras*, Trans. Amer. Math. Soc. **348** (1996), 1191-1212.

[12] R. Megginson, *An Introduction to Banach space Theory*, Springer-Verlag, New York, 1998.

[13] M. Neufang, *Solution to a conjecture by Hofmeier-Wittstock*, Journal of Functional Analysis. **217** (2004), 171-180.

[14] M. Neufang, *On a conjecture by Ghahramani-Lau and related problem concerning topological center*, Journal of Functional Analysis. **224** (2005), 217-229.

[15] J.S. Pym, *The convolution of functionals on spaces of bounded functions*, Proc. London Math Soc. **15** (1965), 84-104.

[16] A. Saghafy, *On the L^1-algebras of some compact totally ordered spaces*, Math. Proc. Camb. Philos. Soc. **122** (1997), 173-184.

[17] A. Ulger, *Arens regularity sometimes implies the RNP*, Pacific Journal of Math. **143** (1990), 377-399.

[18] A. Ulger, *Arens regularity of weakly sequentialy compact Banach algebras*, Proc. Amer. Math. Soc. **127** (11) (1999), 3221-3227.

[19] P.K. Wong, *The second conjugate algebras of Banach algebras*, J. Math. Sci. **17** (1) (1994), 15-18.

Department of Mathematic, Amirkabir University of Technology, Tehran, Iran
Email address: haghnejad@aut.ac.ir

Department of Mathematic, Amirkabir University of Technology, Tehran, Iran
Email address: riazi@aut.ac.ir